


Mark scheme

Question			Answer/Indicative content	Marks	Guidance
1	a		$F (= EQ) = 0.90 \times 1.60 \times 10^{-19} = 1.4(4) \times 10^{-19} \text{ (N)}$	B1	Working and answer must both be shown Answer must be given to 2sf or more Unit need not be given but, if given, must be correct <u>Examiner's Comments</u> This was an easy introduction to the question, which used the definition of electric field strength; $E = F_E / Q$. Being a 'show that' question, candidates needed to show their working in full, including writing the value for the electronic charge (rather than simply 'e') and giving the answer to at least 2 s.f.
	b	i	$(F = BQv \text{ but } B \text{ and } Q \text{ are constant, so})$ (the magnitude of) the velocity is different /changes	B1	Allow speed Ignore the direction is different <u>Examiner's Comments</u> The force on a charged particle moving at right angles to a magnetic field is given by the formula $F_{mag} = BQv$. Since B and Q are constants in this case, the reason for the different magnitude of F must be that the proton has a different velocity, v. Common problems in 6(b)(i) - using the formula $F = BI/\sin\theta$ and suggesting that the proton might be travelling at a different angle to the field, not realising that the proton is always travelling at right angles to the magnetic field in this question - suggesting that the proton may be in a weaker (or stronger) field at X than at Y, not realising that the magnetic field is uniform and so its field strength is constant throughout

			$v = \left(\frac{F_{mag}}{BQ} \right) = \frac{5.6 \times 10^{-19}}{5.0 \times 10^{-5} \times 1.60 \times 10^{-19}}$ resultant force $F_R = (5.6 - 1.4) \times 10^{-19}$ $r = \left(\frac{mv^2}{F_R} \right) = \frac{1.673 \times 10^{-27} \times (7.0 \times 10^4)^2}{4.2 \times 10^{-19}}$ $r = 20 \text{ (m)}$ ii Alternative all-in-one method: $r = \frac{mF_{mag}^2}{F_R B^2 Q^2}$ resultant force $F_R = (5.6 - 1.4) \times 10^{-19}$ $r = \frac{1.673 \times 10^{-27} \times (5.6 \times 10^{-19})^2}{4.2 \times 10^{-19} \times (5.0 \times 10^{-5})^2 \times (1.60 \times 10^{-19})^2}$ $r = 20 \text{ (m)}$	$v = 7.0 \times 10^4 \text{ (m s}^{-1}\text{)}$ implies first C1 Allow 10^{-19} for 1.4×10^{-19} (giving $F_R = 4.6 \times 10^{-19}$) $F_R = 4.2 \times 10^{-19}$ implies second C1 Do not credit if used as F_{mag} in $F_{mag} = BQv$ Third C1 is for correct substitution into formula Allow $m_p = 1.67 \times 10^{-27} \text{ kg}$ given to 3 s.f. Not $m_p = 1.661 \times 10^{-27} \text{ kg}$ or $m_p = 1.675 \times 10^{-27} \text{ kg}$ Allow ECF for incorrect v Use of $F_R = 5.6 \times 10^{-19}$ or $= 1.4 \times 10^{-19}$ is XP Allow $r = 19 \text{ (m)}$ $F_R = 4.2 \times 10^{-19}$ (4.16×10^{-19} to 3sf) implies second C1 An incorrect value of F_R is XP from this point C1 C1 C1 A1 (C1) (C1) (C1) (A1) Third C1 is for correct substitution into formula Allow $r = 19 \text{ (m)}$ <u>Examiner's Comments</u> This question could not be done in one step, by equating the magnetic force to the centripetal force. This is because, at X, the centripetal force is being provided by a combination of forces from both the electric and the magnetic field. The easiest approach is to find the velocity of the proton using $F_{mag} = BQv$ (the value for F_{mag} is given in the diagram as $5.6 \times 10^{-19} \text{ N}$). This velocity v can then be used in the formula $F = mv^2/r$ in order to calculate the radius, r. F here is the <i>resultant</i> force towards the centre of the circle, which is found from magnetic force downwards - electric force upwards (the electric force having been calculated in part (a)). Exemplar 3 is an example of a correct
--	--	--	--	---

				answer, clearly written to show each stage in the calculation: Exemplar 3  The handwritten work shows a diagram with a vertical arrow pointing up labeled 1.44×10^{-19} and a vertical arrow pointing down labeled 5.6×10^{-19}. Below the diagram, the student calculates the resultant force using Pythagoras's Theorem: $F = \sqrt{E^2 + B^2 v^2}$, $F = \sqrt{(1.6 \times 10^{-19})^2 + (5 \times 10^{-19})^2}$, $F = 5.2 \times 10^{-19}$. They also calculate the radius: $r = \frac{mv^2}{F}$, $r = \frac{(1.67 \times 10^{-27}) \times (5000)^2}{5.2 \times 10^{-19}}$, $r = 8.1 \times 10^{-5}$ m.
		iii	$\text{resultant force} = (\sqrt{3.9^2 + 1.4^2}) \times 10^{-19}$ $\text{resultant force} = 4.1 \times 10^{-19} \text{ (N)}$	Ignore attempt to calculate weight of proton Allow $F_E = 10^{-19}$ Allow $F = 4.0 \times 10^{-19} \text{ (N)}$ using $F_E = 1.0 \times 10^{-19}$ Allow $F = 4.2 \times 10^{-19} \text{ (N)}$ using $F_E = 1.44 \times 10^{-19}$ Examiner's Comments There are two forces acting on the proton at Y: an electric force upwards (given in (a)) and a magnetic force to the left (shown on the diagram). These two forces act at right angles to each other, and so the magnitude of their resultant can be found using Pythagoras's Theorem. Credit was given for using a value for the electric force to 1, 2 or more significant figures.
		iv	<u>resultant / net</u> force is not perpendicular to velocity work is done on proton (therefore kinetic energy changes so speed is not constant)	Allow direction of motion / path but not speed for velocity Allow acceleration / <u>resultant</u> force is not (always) towards centre (of circle) Allow electric force is not perpendicular to velocity / is in the same direction as velocity Ignore references to centripetal Ignore references to centripetal Examiner's Comments At Y, the proton is moving downwards, with a resultant force being the combination of an electric force upwards and a magnetic force to the left (calculated in part 1). The resultant force cannot be at right angles to the

					velocity, so we cannot have circular motion. The component of the resultant force acting in the direction of the proton's motion will do work on the proton and change its speed. So, the proton cannot be travelling at a constant speed.
			Total	10	
2	a	i	$\frac{GMm}{r^2} = \frac{mv^2}{r}$ $\frac{1}{2}mv^2 = \frac{1}{2} \frac{GMm}{r}$	M1 A1	Allow omission of 'm' on both sides of equation (gravitational field strength = centripetal acceleration) Cancelling/rearrangement/identification of GMm/r as GPE <u>Examiner's Comments</u> Many candidates made a good start by equating the formula for gravitational force with the expression for centripetal motion. Others that assumed that g of 9.81 m s^{-2} did not score any marks. The simplest way to arrive at the correct expression was to identify GM/r in the gravitational force formula, to rearrange and then multiply both sides by $\frac{1}{2}$. Approximately $\frac{1}{2}$ of all candidates that responded got as far as this gaining both marks.
		ii	(Increase in) GPE = $(-56 - -63 \text{ MJ}) = 7 \text{ (MJ)}$ or (Increase in) KE = $0.5 \times 56 = 28 \text{ (MJ)}$ Sensible reasoning, e.g. $7+28 > 30$	M1 A1	Allow evaluation of total energy of 35 (MJ) <u>Examiner's Comments</u> Many candidates correctly determined how much GPE the satellite needed to gain i.e. 7 MJ in order to reach orbit from -63 MJ to -56 MJ. To find the KE when in orbit, candidates needed to use the result from the previous part of the question. This explains why the in orbit, the KE required is $\frac{1}{2}$ of 56 MJ i.e. 28 MJ. A small fraction of candidates successfully accomplished this step.

					This means the total energy gain is the sum of 28 MJ and 7 MJ i.e. 35 MJ.
	b		Level 3 (5–6 marks) Correct calculations, and advantages and limitations discussed. <i>There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated.</i> Level 2 (3–4 marks) Correct calculations and an advantages discussed or Correct calculations and a limitation discussed. <i>There is a line of reasoning presented with some structure. The information presented is in the most part relevant and supported by some evidence.</i> Level 1 (1–2 marks) Attempted calculations or a single correct calculation or incomplete explanations of advantages and/or limitations. <i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant.</i> 0 mark <i>No response or no response worthy of credit.</i>	B1 × 6	Use level of response annotations in RM Assessor Ignore general knowledge answers e.g. accidents, cost, politics Allow references to energy, energy per unit mass or potential as interchangeable Allow ecf from candidate's value for total energy per unit mass in orbit from (i.e. 35 MJ) 22bii or use of 30 (MJ) Indicative scientific points may include: Calculations • (Minimum) additional KE from aircraft=26 kJ • Additional GPE from aircraft = 100 kJ • Additional KE from equatorial launch=110 kJ • GPE calculated by mgh as an acceptable approximation • (Without taking Earth's rotation into account,)KE at equator is about 4x aircraft KE • GPE calculated from $vg = (-)GM/r$ • Total energy = $0.5GM/r$ Advantages • Aircraft launch provides KE and GPE • Aircraft velocity will be higher than 230 depending on where the aircraft takes off. • Less (rocket) fuel required • Aircraft launch has similar/slightly larger energy to equatorial • Equatorial launches can only happen in limited locations/aircraft launches can take place almost anywhere Limitations

				• Aircraft launches only suitable for small satellites. • Effects (for either) only significant for near earth orbits/low altitudes • Either launch provides very small fraction (less than 1%) of the energy required Examiner's Comments Many candidates found this difficult to access. A solely qualitative evaluation was limited to level 1 (2 marks). Exemplar 2 <i>The initial kinetic energy + GPE will give the total energy gained by the satellite. On earth $\frac{1}{2} \times 1 \times 400^2 = 0.1058 \text{ MJ}$ of kinetic energy. On the aircraft, $\frac{1}{2} \times 1 \times 230^2 = 0.02645 \text{ MJ}$. However, on the aircraft, the GPE will be higher, and the satellite will need to gain less GPE. $\frac{6.07 \times 10^6 \times 6 \times 10^{-3}}{6.37 \times 10^6} = \text{GPE advantage.}$ Earth = 10000</i> <i>In conclusion, although the aircraft would slightly reduce energy required for GPE increase, it would reduce initial kinetic energy by a factor of 4, and there would also be energy required to power the aircraft.</i> In Exemplar 2, the candidate has completed a small number of calculations comparing the KE the two methods would raise. There is also a statement that these differ by a factor of 4 along with an attempt at calculating the GPE advantage by launching from the aircraft rather than from the ground. Crucially, there is very little mention of limitation and no supporting calculations. Holistically speaking, therefore, there was not enough evidence to award this candidate Level 3, yet sufficient for a Level 2. Exemplar 3
--	--	--	--	---

					which starts with maximum gradient and then flattens out.
		ii	Resistive force increases (with speed) Zero net or zero resultant force	B1 B1	Allow drag / (air) resistance / friction for 'resistive force' Allow resistive force = component of weight down the slope NOT simply idea of resistive force = weight <u>Examiner's Comments</u> While many candidates appreciated that the car reached a maximum speed because the resultant force was zero, some contradicted this by saying that the weight = drag (as it would be in vertical motion) or something else incorrect. Far fewer candidates stated that the drag increases with speed effectively. Quoting the given expression $F = kv^2$ was deemed insufficient. Examination Tip Repeating information given in the question is rarely creditworthy by itself.
		iii	Component of weight down slope = $9300 \sin 5^\circ$ Re-arrange to $(k=)F \div v^2$ $(k=)810 \div 900 = 0.9\dots$	M1 M1 A1	Allow 810 or 811 seen Allow substitutions for variables Mark is for substitution <u>and</u> candidate's value seen <u>Examiner's Comments</u> As this question is a 'show that', all steps were required. Many candidates omitted the rearrangement stage, restricting their maximum score for this item to 1 mark. This approach was consistent throughout the paper for this type of question. Examination Tip Make sure that all steps of working are presented in 'show that' questions, especially the step that shows the relevant quantity as the subject of the equation. Always show your evaluation

					to at least 1 more significant figure than that shown in the question.
	b		evidence of substitution of $F=kv^2$ into $P = Fv$ $v = (P \div k)^{1/3}$ $v = 44 \text{ (m s}^{-1}\text{)}$	C1 C1 A1	e.g. $P = kv^3$, $P = (kv^2) v$, etc Allow use of $k = 1$ which gives 42 Allow answer within range 36 to 53 <u>Examiner's Comments</u> The key idea here is that the force from the engine (given by $F = P / v$) will equal the resistive forces ($F = kv^2$) when the car is at maximum speed. Candidates could choose which value of k they used here, either $k = 1$ from the question data or the value of k from the previous item. This gives an acceptable range of speeds as stated in the mark scheme.
	c		Power is proportional to the speed cubed / Max speed is proportional to the cube root of max power / power proportional to speed $\times kv^2$ Valid reference to the cube root of 2 increase in velocity for double power / Valid reference to factor of 8 increase in power for double the velocity	B1 B1	NB cube root of 2 is 1.2599... e.g. $1.26 \times 44 = 55 \text{ (m s}^{-1}\text{)}$ <u>Examiner's Comments</u> Even if they couldn't complete the calculation in the previous item, candidates needed to be able to state the idea qualitatively for the first mark. No further calculations were required, except the correct answer that the maximum speed would increase by a factor of cube root (2).
			Total	12	
4		i	Horizontal = $43 \text{ (.3) (m s}^{-1}\text{)}$ Vertical = $25 \text{ (m s}^{-1}\text{)}$	B1 B1	ignore sign Allow answers in incorrect order for 1 mark MAX Allow $H = 7.7 \text{ (m s}^{-1}\text{)}$, $V = -49.4 \text{ (m s}^{-1}\text{)}$ 1 mark MAX <u>Examiner's Comments</u> Most candidates got this correct. Some scored a single mark for confusing sine and cosine but otherwise completing the calculation correctly. Examination Tip Make sure to check whether your calculator should be in degrees or

					radians mode. Also, practise which trigonometric function to use. If in doubt, draw a labelled right-angled triangle to help identify the correct function.
		ii	Correct application of N3L Plus ONE from: (Direction of) momentum of air has changed or direction of air flow has changed There is a force on the air (from the model)	B1 B1 B1	force on air is equal (and opposite) to force on model <u>Examiner's Comments</u> The key idea here was Newton's third law, with credit for some supporting information.
		iii	$F = \frac{\Delta p}{\Delta t}$ $F = 35 \times 25$ (divided by 1) $F = 880$ (N)	C1 C1 A1	ecf candidate's vertical velocity in (i) Allow 875 Allow 870 <u>Examiner's Comments</u> Answers that clearly used $F = ma$ with the acceleration of 25 ms^{-2} were deemed wrong Physics (as in previous series). Error Carried Forward (ECF) was clearly applied here also.
			Total	7	
5			Similarity: same magnitude or same speed or (still) $0.17 \text{ (m s}^{-1}\text{)}$ Difference: (different) direction or (approximately) opposite direction	B1 B1	Allow ECF from (a) for 0.17 Not same velocity Allow negative <u>Examiner's Comments</u> Most candidates understood that velocity was a vector quantity and there were many correct answers explaining the similarity in that the magnitude of the velocity was the same and the difference was the direction. Lower scoring candidates often stated that the velocity was the same. This suggests that there was not the full understanding of the physical quantities speed and velocity.
			Total	2	

6	a	$22.3 \cos 84 (= 2.33)$ 2.33	M1 A0	ALLOW $22.3 \sin 6$ <u>Examiner's Comments</u> Most candidates clearly showed how the initial velocity of the ball could be resolved into its horizontal component. Most candidates used $\cos 84^\circ$ although there was a significant minority who correctly used $\sin 6^\circ$.
	b	$v_v^2 = 2.33^2 + 2 \times 9.81 \times 2.4$ $v_v = \sqrt{52.517} \text{ OR } 7.247$ 7.25	M1 M1 A0	ALLOW ECF from (a) <u>Examiner's Comments</u> For this part of the question it was necessary for candidates to clearly show the substitution of the data into the equation. This includes the value of g (as 9.81 m s^{-2}) from the data sheet. It was also necessary for candidates to show that having determined v^2, this value needed to have the square root taken. Higher performing candidates wrote the final answer as 7.247 or $7.24685 \approx 7.25 \text{ (m s}^{-1}\text{)}$.
	c	$u_h = 22.3 \sin 84 = 22.2$ $v = \sqrt{22.2^2 + 7.25^2} = 23.35$ $(= \frac{1}{2} \times 0.210 \times 23.35^2) = 57.2 \text{ (J)}$ OR Change in potential energy = $0.210 \times 9.81 \times 2.40 = 4.94$ Initial kinetic energy = $\frac{1}{2} \times 0.21 \times 22.3^2 \text{ OR } 52.2 \text{ (J)}$ $(4.94 + 52.2 =) 57.1 \text{ (J)}$	C1 C1 A1 C1 C1 A1	ALLOW $v^2 = 545$ ALLOW 57.2 (J) <u>Examiner's Comments</u> This question proved challenging to candidates. The common error was to calculate the kinetic energy using a value of 7.25 m s^{-1}. There were two main methods of answering this question: Either: determining the horizontal component of the velocity of the ball, (which remains constant) then working out the resultant velocity of the ball as it hits the ground and then calculate the kinetic energy. Or: Calculate the change in gravitational

					potential energy Calculate the initial kinetic energy of the ball And then add the two values together.  Misconception Omitting the kinetic energy in the horizontal direction when calculating the kinetic energy of the ball. Candidates should be able to compare the motion of a projectile in two perpendicular directions and also confirm that similar results are obtained by considering energy transfers.
	d		(change of) Momentum = mass ×(change of) velocity As velocity increases, the momentum increases OR force = rate of change of momentum gravitational force acts on the ball and increases momentum OR Momentum is a vector quantity, change in direction means that the momentum changes.	M1 A1 M1 A1 M1 A1	ALLOW changes for increases <u>Examiner's Comments</u> There are many possible explanations as to why the momentum of the ball changes. To score full marks candidates needed to state a property of momentum, e.g. momentum = mass × velocity before then explaining why the momentum would change, e.g. as the ball falls, velocity increases so for constant mass the momentum increases.
			Total	8	
7		i	newton in base units is kg m s^{-2} Substitution and cancelling of kg and m arriving at $\text{s}^2 \rightarrow \text{s}^2$	C1 A1	
		ii	One force is increased by kx and one is reduced by kx /AW Some working to include $kx - (-kx)$	B1 B1	reject 2 springs in series or 2 springs in parallel idea XP accept one extension is reduced by x and one is increased by x / AW <u>Examiner's Comments</u> Question 21 (a) (ii) is considerably more challenging. The two springs are not in series nor are they in parallel.

					When there is a displacement x one spring is extended by an <i>extra</i> amount x i.e. an extension of $(e + x)$ and the other is extended by a reduced amount x i.e. an extension of $(e - x)$ where e is the equilibrium extension. This meant that the resultant force was $k(e + x) - k(e - x)$, which is clearly $2kx$. Neither spring goes into compression, although we condoned candidates who suggested that a reduction in extension meant the same as a compression.
		iii	period is independent of <u>amplitude</u> / AW No effect	M1 A1	Allow isochronous <u>Examiner's Comments</u> A reasonably large proportion of candidates did not link the idea of initial displacement to the amplitude of this motion. Those that did often scored both marks as they also recalled that SHM is isochronous.  Assessment for learning Merely repeating the words in the question, in this case 'the initial displacement' instead of 'amplitude', is unlikely to give access to full marks. Think about which piece or pieces of technical language on the specification are the likely target of each question.
			Total	6	
8	a	i	Arrow along the line of the support rod labelled tension or T.	B1	Allow unlabelled single arrow along either rod Allow unlabelled arrows along both rods Allow arrow(s) up, down or both NOT any contradictory arrows <u>Examiner's Comments</u> In Question 17 (a) (i) of this question, the phrase 'tension in the rod' can mean several different things, all of which were given in the mark scheme.

		ii	11.1 sin 35 or 11.1 cos 55 seen addition of 3.9 (half the diameter of the support disc) to candidate's horizontal component of rod length Total= 10.3m	M1 M1 A0	NOT use of tan 35 or tan 55 allow 7.8/2 for 3.9 10.27 to 2 dp NB use of 11.1 cos 35 or 11.1 sin 55 arriving at 12.99 scores 1 (wrong trig) NB reject use of radians (scores 0) <u>Examiner's Comments</u> Many candidates approached part (ii) with some confidence, spotting that the horizontal portion of the rod was 11.1 sin(35) and that it should be added to the radius of the disc.
		iii	$mg = T \cos 35$ $T = mg \div \cos 35$ = 140 N	C1 C1 A1	Allow use of sin 55 NOT use of tan 35 or tan 55 Answer is 143.7 N to 4 sf <u>Examiner's Comments</u> Parts (a) (iii) and (iv) were more challenging, requiring good knowledge of both circular motion and how to calculate components of forces. Again there were several legitimate routes to the right answer, all of which were mentioned in the mark scheme. Very logical approaches were in part (a) (iii), to equate the vertical component of the tension with the weight of the sandbag.
		iv	$T \sin 35 = mr\omega^2$ $\omega = \sqrt{\frac{T \sin 35}{mr}}$ =0.8(17) radian s⁻¹	M1 A1 A0	Allow use of $W \tan 35$ or $W \tan 55$ Allow use of cos 55 and/or mv^2/r Allow use of Pythagoras to find centripetal force (82.4...) NOT use of $T \tan 35$ or $T \tan 55$ Allow ω^2 subject. Allow any combination of rearrangement and substitution ECF allowed for T and r. Use of 2 s.f. values for T and r gives 0.84 m <u>Examiner's Comments</u>

					In part (a) (iv), the quickest approach was to equate the horizontal component of the tension with the centripetal force. The data booklet provides a convenient expression for the centripetal force in terms of the angular velocity, without the need for finding the tangential velocity.
	b	i	Use of $17 = \frac{1}{2}gt^2$ $= 1.9 \text{ (1.86) s}$	C1 A1	i.e. substitution of 17 and g or 9.81 or 9.8 e.g. $s = (ut) + \frac{1}{2}at^2$ $t = \sqrt{\frac{2s}{a}}$ $= \sqrt{\frac{2 \times 17}{9.81}}$ Allow any subject
		ii	Horizontal speed = ωr or Horizontal distance = speed $\times$ time $= 0.82 \text{ radians s}^{-1} \times 10.26 \text{ m} \times 1.86 \text{ s}$ $= 16 \text{ m (15.6 m)}$	C1 C1 A1	Use of data in the question stem (0.8 and 10) allowed, which gives 15.2 m. Ecf for use of candidate's value of r and ω, giving values rounding to between 14.9 and 16.0 m
		iii	Relevant variable identified Effect on speed of shoe or time of flight of shoe correctly identified Conclusion consistent with relevant physics e.g. - Shoe is lower mass yet no change in angular velocity or radius since independent of mass so no change in horizontal displacement. - Shoe is below seat so would be travelling with larger radius/speed so larger distance travelled horizontally - Shoe might have been kicked off backwards so have lower speed so lower distance	M1 M1 A1	e.g. mass/weight, drag/air resistance, radius, height, starting condition (e.g. kicking shoe off) Assume "it" in response refers to the shoe. ignore velocity for first M1 allow correct explanation of "no effect" on speed or time by change of mass <u>Examiner's Comments</u> Question 17 (b) explored ideas about parabolic flight due to gravitation. The only force acting on the sandbag and the shoe after they have been released is the weight force, which acts vertically downwards. Many candidates realised that the vertical velocity of the sandbag when it left the swing was zero, enabling them to calculate the time for the bag to fall 17m vertically downwards (using $s = \frac{1}{2}gt^2$)

			- Shoe would come from below the seat/lower than the sandbag i.e. vertical distance to fall less, thus time of flight and horizontal distance less. - Effect of air resistance hadn't been included so shoe suffers drag, decelerating horizontally so distance would be smaller		To calculate the horizontal distance travelled required both the horizontal velocity (from $v = r\omega$) and the time of flight from part (b) (iii). There was lots of scope for applying error carried forward rules as mentioned in the mark scheme.  Misconception In lots of questions, candidates make assumptions when trying to use formulae to justify their ideas. In this case, it was that for the same radius, the shoe must leave the seat faster than the sandbag, purely because the shoe had less mass. Often, what is constant is as important to consider as what is changing. Here, if the radius for the sandbag and the shoe are the same, then the horizontal velocity at release must be the same, since the radius and angular speed for both are the same, using $v = r\omega$.
			Total	16	
9			C	1	
			Total	1	
10			B	1	<u>Examiner's Comments</u> This question was generally answered well as most candidates correctly resolved T, tension to determine an expression for W.
			Total	1	
11			C	1	<u>Examiner's Comments</u> The first step with this question is to calculate the resultant force, which is 0.2 N upwards (eliminating option A). As the drag force is upwards, the direction of motion must be downwards (eliminating option D). Since the

					resultant is opposite to the direction of travel, this object must be decelerating.
			Total	1	
12			B	1	
			Total	1	
13			D	1	<u>Examiner's Comments</u> Most candidates got this correct by correctly identifying mass as a scalar and force as a vector. The most likely incorrect answer was C.
			Total	1	
14		i	$a = \frac{6.0}{3.0}$ 2.0	C1 A0	Allow any correct gradient calculation <u>Examiner's Comments</u> Candidates had to show that the acceleration was 2.0 m s^{-2} which 90% of candidates demonstrated successfully by using values from the graph and calculating a gradient value which equalled the acceleration. To be given this mark, candidates had to clearly show their working out using values taken from the graph.
		ii	$680\cos 55 / 150 \times 2.0$ $680\cos 55 - R = 150 \times 2.0$ $R = 90 \text{ (N)}$	C1 C1 A1	If both components given (vertical and horizontal) it must be clear that the 390N is the horizontal component. <u>Examiner's Comments</u> Candidates performed less well on this question as it was mostly only the most successful responses that were given 3 marks for resolving the horizontal component of the tension in the rope to correctly calculate the horizontal resistance R. About 40% of candidates were given 1 mark for either correctly resolving the horizontal component of the tension to give 390 N or calculating the resultant force ($F = ma$) to give 300 N. It would have helped candidates understand the question to draw a free

					body force diagram to identify the forces acting on the buggy and direction and magnitude of the resultant force.
			Total	4	
15	a	i	$E_p (= 0.16 \times 9.81 \times 2.5) = 3.9 \text{ (J)}$	A1	3.924 <u>Examiner's Comments</u> This question was generally answered well.
		ii	$v^2 = \frac{2 \times 3.9}{0.16}$ or 48.75 OR $v^2 = 2 \times 9.81 \times 2.5$ or 49.05 $v = 7.0 \text{ (ms}^{-1}\text{)}$	C1 A1	Allow ECF from (a)(i) Allow 1sf <u>Examiner's Comments</u> Most candidates correctly equated the change in gravitational potential energy to kinetic energy and gained the correct answer. Other candidates correctly used $v^2 = 2 gh$. A number of candidates gave the answer as 7 (ms ⁻¹) – since the data in the question was given to two significant figures the answers should also be given to two significant figures. Ideally the substitution of data into appropriate equations should be shown.
	b	i	$R (= 12 \times 0.71) = 8.5(2) \text{ (m)}$	A1	<u>Examiner's Comments</u> It was expected that candidates would multiply the horizontal velocity by the time. This was generally answered well.  Assessment for learning When considering projectile motion, candidates should treat the vertical and horizontal velocities independently.
		ii	$E_k = \frac{1}{2} \times 0.16 \times 12^2$ or 11.5 OR $= \frac{1}{2} \times 0.16 \times 13.9^2$ or $\frac{1}{2} \times 0.16 \times 193$	C1	Allow use of vertical $v = 6.97$ (calculated using $v = u + at$;))

			$E_k (= 11.5 + (a)(i) = 11.5 + 3.9 = 15(.4) \text{ (J)}$	A1 Allow 15.5 (J) Allow ECF from (a)(i) <u>Examiner's Comments</u> Many candidates calculated the kinetic energy of the ball using the velocity of the ball in the horizontal direction but then did not add the change in potential energy of the ball as it fell. Other candidates determined the resultant velocity of the ball and then calculated the kinetic energy.
		iii	$\theta \left(= \tan^{-1} \left(\frac{(a)(ii)}{12} \right) = \tan^{-1} \left(\frac{7}{12} \right) \right) = 30^\circ$	A1 Allow ECF from (a)(ii) 30.256 <u>Examiner's Comments</u> Candidates achieving on this question correctly determined the angle using the horizontal and vertical velocities. Where the response was incorrect, candidates had used either energies or distances.
			Total	7